



Study The Effect of Different Phosphorus Fertilization Resources on The Growth and Productivity of German-Chamomile (*Matricaria chamomilla* L.) Plant under Sand Soil Conditions

Wafaa Hamed Abd El-Alem

Desert Research Center, Medicinal and Aromatic Plants Department, 11753 Cairo, Egypt

Received: 11 June 2022

Accepted: 12 July 2022

Published: 30 July 2022

ABSTRACT

This research was conducted at Sekem farm in the desert back of Minya Governorate, Egypt, during the two consecutive seasons of 2018/2019 and 2019/2020. The work aimed to study the effect of different phosphorus fertilization resources on the growth, yield, and essential oil content of chamomile (*Matricaria chamomilla* L.). The treatments included six phosphorus fertilization resources, like the following: F1: control 48 kg P₂O₅ (single super phosphate 16% P₂O₅), F2: 48 kg P₂O₅ (phosphoric acid 85% P₂O₅), F3: 48 kg P₂O₅ (24 kg P₂O₅ single super phosphate 16% P₂O₅ + 24 kg P₂O₅ phosphoric acid 85% P₂O₅), F4: 48 kg P₂O₅ (mono ammonium phosphate MAP 12% N:61% P₂O₅), F5: 48 kg P₂O₅ (24 kg P₂O₅ single super phosphate 16% P₂O₅ + 24 kg P₂O₅ mono ammonium phosphate), and F6: 48 kg P₂O₅ (16 kg P₂O₅ single super phosphate 16% P₂O₅ + 16 kg P₂O₅ phosphoric acid 85% P₂O₅ + 16 kg P₂O₅ mono ammonium phosphate MAP 12% N:61% P₂O₅). Results showed that applying 48 kg mono ammonium phosphate (MAP 12% N: 61% P₂O₅) produced the highest plant growth parameters, oil content, N, P, and K in plant tissues. Also, this treatment increased the concentrations of chamazulene to 41.68 % and bisabolol oxide-B to 13.61%. In comparison, as traditional phosphorus fertilizer, super-phosphate gave the lowest values in the two tested seasons.

Keywords: *Matricaria chamomilla* L., phosphorus fertilization, yield, chamazulene.

1. Introduction

Medicinal and aromatic plants are of fundamental economic importance because of local and foreign markets' continuous and increasing demand for their products (Khalid, 2006; Hamed, 2018; Abd El-Alem *et al.*, 2021; Toaima, 2022). Chamomile is a common name for several daisy-like plants of the family Asteraceae (Presibella *et al.*, 2006). The vital kind of chamomile is German chamomile (*Matricaria chamomilla* L.). Its native region is southern and eastern Europe, where it spreads. The economic segment: inflorescences being dried and essential oil. Egyptian chamomile is in great demand for foreign markets. Fayoum, Beni Suef, Minya, and Assiut are the most productive areas. The average productivity per feddan is about 500-600 kg of entire inflorescences and gives about 3-8 kg volatile oil in the season. The inflorescences contain volatile oil ranging between 0.70 - 0.77%, extracted by steam distillation. The resulting oil is distinguished by its blue color because of its chamazulene presence, which has a 38.6 - 41.8% ratio. Chamazulene determines the quality of the produced oil. Also, the oil contains other substances such as bisabolol oxide B, bisabolol, farnesene, germacrene, and bisabolol oxide A (Das, 2014). Chamomile is used to treat indigestion cases, irritable bowel syndrome, general tonic, antipyretic, soothing nerves, and treat insomnia, encouraging stomach and bile secretions. Treating skin infections and wounds, making anti-fever medications, anti-colic and convulsions, gain on flavor in drinks, dental pain, mouth and gum ulceration, gout treatment, anti-inflammatory, anti-allergic, anti-cramping, eye fatigue and fatigue, manufacture of cosmetics, soaps, and perfumes (Sheibanivaziri, 1997; Aba *et al.*, 2004; Omidbeigi, 2005; Salmon and Abou-zied, 2006; Letchamo *et al.*, 2006; Ahmadian *et al.*, 2011).

Phosphorus is a moving component within the plant that is a slow movement in the soil, and it is one of the fundamental nutrients in plant nutrition. It comes second after nitrogen in terms of its quantity

Corresponding Author: Wafaa Hamed Abd El-Alem, Desert Research Center, Medicinal and Aromatic Plants Department, 11753 Cairo, Egypt. E-mail: dr.wafaa_60@yahoo.com

in plant tissues. Microorganisms transform organic phosphorous into inorganic phosphorous. The degree of phosphorous utilization of a plant varies according to several factors. The most important factors are:- the type of clay metal, fixed in clay and sandy lands with a higher calcium content than light and pH in the soil where the highest validity level is to benefit from it at pH 6.5-7. The CO₂ gas released from the decomposition of organic matter plays a significant role in increasing the susceptibility of phosphorus to benefit. The essential functions of phosphorus are included in the composition of the nucleus protein, which is a necessary component of respiration processes. It has a role in transforming carbohydrates inside the plant, such as converting starch into sugar. It has a role in representing fats and accelerates the fruit ripening processes. Abla *et al.*, (2004) reported that phosphorus had a pronounced effect on carbohydrates, soluble sugars, mineral contents, and the percentage of oil production from chamomile flowers compared with the control.

The problem is that calcium superphosphate quickly becomes a non-absorbable form in the soil after a short addition period. The symptoms of P element deficiency appear especially with lower temperatures and the entry of plants into the flowering stage (Saharkhiz and Omidbaigi, 2008). The most commonly used fertilizer phosphorus sources are phosphoric acid and mono ammonium phosphate (MAP). These phosphorus fertilizer sources are highly water-soluble (>80 %) and thus dissolve quickly in soil to release plant-available phosphate and ammonium. Now MAP and phosphoric acid are widely used as sources of P fertilizer. They are popular in most European countries due to their high P nutrient content and excellent physical properties (Rosen *et al.*, 2014).

This research aims to study the effect of the different phosphorous sources that are more readily available for absorption on the growth, yield, and active ingredients of the chamomile plant instead of the traditional forms such as calcium superphosphate. Also, to increase the productivity of chamomile plants under conditions of reclaimed lands, most of which are exposed to frost waves for long periods that affect productivity in Egyptian desert lands (sandy soils).

2. Materials and Methods

The field experiment was conducted at Sekem farm in the desert back of Minya Governorate, Egypt, during the two consecutive seasons of 2018/2019 and 2019/2020. The aim was to study the effect of different phosphorus fertilizers on the growth, flower yield components, and volatile oil production of chamomile (*Matricaria chamomilla*) plants.

Chamomile seeds were obtained from the National Research Centre, Medicinal and Aromatic Plants Department, Dokky, Giza, Egypt. Seeds were sown on 25th and 30th September during the 1st and 2nd seasons, respectively. The sowing was directly in the reclaimed sandy soil plains. 15 cm between the lines and 7 cm between the plants using a seed planting machine at a rate of 1 kg seed per feddan. The seeds index 0.1 g (100-120 plant / m²).

Plants were irrigated by central spraying at a rate of 20-25 m³ per feddan until germination is complete, then 20 m³ per day after day. 90 kg of nitrogen was added as ammonia sulfate + 100 kg of potassium sulfate per feddan. It was added injection in the irrigation system throughout the season + 5 tons of compost / feddan, added to all treatments during soil preparation for cultivation (Sharafzadeh *et al.*, 2011; Naderidarbaghshahi *et al.*, 2011; Osman *et al.*, 2013). The physical and chemical properties of the soil are shown in Table (1). The analyses were conducted according to Chapman and Pratt (1978). Also, the chemical properties of irrigation water and compost manure are shown in Tables (2 and 3).

Table 1: The mechanical and chemical analyses of the experimental soil area

Depth (cm)	Sand (%)	Silt (%)	Clay (%)	Soil texture						
0-30	86.00	5.00	9.00	Sandy						
pH	E.C. (ds/m)	O.M. (%)	Soluble anions (meq/l ⁻¹)				Soluble cations (meq/l ⁻¹)			
			CO ₃ ²⁻	HCO ₃ ⁻	Cl ⁻	SO ₄ ²⁻	Ca ⁺⁺	Mg ⁺⁺	Na ⁺	K ⁺
7.90	1.80	0.5	-	1.00	10.00	7.00	6.00	8.00	8.60	1.40

Table 2: The chemical analysis of irrigation water

pH	E.C.	Soluble anions	Soluble cations
----	------	----------------	-----------------

	(ppm)	(meq/l ⁻¹)				(meq/l ⁻¹)			
		CO ₃ ⁻	HCO ₃ ⁻	Cl ⁻	SO ₄ ⁻	Ca ⁺⁺	Mg ⁺⁺	Na ⁺	K ⁺
7.32	747.15	-	4.26	8.59	7.44	11.45	5.64	7.31	0.69

Table 3: The chemical analysis of used compost manure

Bulk density kg/m ³	Moisture content (%)	pH	EC (ds/m)	O.M. (%)	Ash (%)	C/N ratio (%)	N (%)	P (%)	K (%)
639	25	7.91	5.53	25.97	74.03	10.91	1.38	0.49	0.59

The treatments included six different phosphorus fertilization resources, like the following:

- F1:** Control 48 kg P₂O₅/feddan (single super phosphate 16% P₂O₅), divided into two batches (30 kg when preparing the soil for sowing, then 18 kg when flowering begins).
- F2:** 48 kg P₂O₅/feddan (phosphoric acid 85% P₂O₅), distributed over 24 weeks from the cultivation date.
- F3:** 48 kg P₂O₅/feddan (24 kg P₂O₅ single super phosphate 16% P₂O₅ + 24 kg P₂O₅ phosphoric acid 85% P₂O₅), distributed over 24 weeks from the start of cultivation.
- F4:** 48 kg P₂O₅/feddan (mono ammonium phosphate MAP 12% N:61% P₂O₅), distributed over 24 weeks from cultivation.
- F5:** 48 kg P₂O₅/feddan (24 kg P₂O₅ single super phosphate 16% P₂O₅ +24 kg P₂O₅ mono ammonium phosphate), distributed over 24 weeks from cultivation.
- F6:** 48 kg P₂O₅/feddan (16 kg P₂O₅ single super phosphate 16% P₂O₅ + 16 kg P₂O₅ phosphoric acid 85% P₂O₅ +16 kg P₂O₅ mono ammonium phosphate MAP 12% N:61% P₂O₅), distributed over 24 weeks from cultivation.

Harvesting was conducted three times during the growing season, on 5th January, 20th February, and 15th April, and samples were dried in the shade (Kandeel, 1982; Essa, 1999). The recorded data included:

2.1. Vegetative growth parameters

Plant fresh weight (g/m²), plant dry weight (g/m²), fresh weight of flower heads (g/m²), and dry weight of flower heads (g/m²).

2.2. Chemical constituents

2.2.1. Essential oil percentage: was determined during both seasons and extracted from the air-dried flower heads by hydrodistillation using a Clevenger apparatus (British Pharmacopoeia, 1963).

2.2.2. Essential oil yield/m² (ml): This was calculated as follows: oil percentage × flower heads dry weight per m²/100.

2.2.3. Essential oil composition

GC-MS analysis of essential oils was performed by Gas Chromatography-Mass Spectrometry instrument at the Laboratory of Medicinal and Aromatic Plants, National Research Center, Egypt, with the following specifications. Device: a TRACE GC Ultra Gas Chromatographs (THERMO Scientific Corp., USA), coupled with a THERMO mass spectrometer detector (ISQ Single Quadrupole Mass Spectrometer). The GC-MS system had a TR-5MS column (30 m x 0.32 mm i.d., 0.25 µm film thickness). Analyses were conducted using helium as carrier gas at a flow rate of 1.3 ml/min at a split ratio of 1:10 and the following temperature program: 80°C for 1 min; rising at 4°C/min to 300°C and held for 1min. The injector and detector were held at 220 and 200°C, respectively. Diluted samples (1:10 hexane, v/v) of 1 µL of the mixtures were continuously injected. Mass spectra were obtained by electron ionization (EI) at 70 eV, using a spectral range of m/z 40-450. Identification of the compounds depended on comparing the retention times with those of authentic samples. Comparing their linear retention indices relative to the series of n-hydrocarbons and on computer matching against commercial and

libraries mass spectra built up from pure substances, components of known oils, and MS literature data (Bunzen *et al.*, 1969; Hoftman, 1967; Massada, 1976; Jennings and Shibamoto, 1980; National Committee for Clinical Laboratory Standards, 2002; Adams, 2007; Babushok *et al.*, 2011).

2.2.4. Mineral elements content

Nitrogen, phosphorus, and potassium percentages were determined according to Koch and McMeekin (1924), Troug and Mayer (1939), and Brown and Lilleland (1946), respectively.

The randomized complete block design was used with three replicates (six treatments). According to Snedecor and Cochran (1968), the statistical analysis was carried out, and L.S.D. test at 5 % was used to compare the means.

3. Results and Discussion

3.1. Vegetative characters

3.1.1. Fresh and dry weights of plant (g/m²)

Data in Table (4) presented fresh and dry weights of plants (g/m²). Results showed that all treatments increased the fresh and dry weights of *Matricaria chamomilla* plants compared with the control treatment. This increment is significant in both seasons. Compared with other fertilization treatments, the best results were obtained with F4 48 kg P₂O₅/feddan (mono ammonium phosphate). This treatment gave the heaviest fresh and dry weights of plants in the 1st and 2nd seasons, respectively which recorded 1510 and 1515 g/m². Also, the same treatment produced the highest value for dry weights of the plant (g/m²), which recorded 345.33 and 350.66 g/m². It was followed by F6 48 kg P₂O₅ (16 kg P₂O₅ single super phosphate 16% P₂O₅ +16 kg P₂O₅ phosphoric acid 85% P₂O₅ +16 kg P₂O₅ mono ammonium phosphate MAP 12% N: 61% P₂O₅). Meanwhile, fertilization with control treatment F1 48 kg P₂O₅ (single super phosphate 16% P₂O₅) produced the lowest fresh and dry weights in the first and second seasons, respectively.

Table 4: Effect of different phosphorus fertilization resources on the plant fresh and dry weights (g/m²) during 2018/2019 and 2019/2020 seasons

Treatments	Plant fresh weight (g/m ²)			Plant dry weight (g/m ²)		
	1 st season	2 nd season	Mean	1 st season	2 nd season	Mean
F1	1350.00	1365.00	1357.50	275.33	276.67	276.00
F2	1410.00	1400.00	1405.00	289.33	288.67	289.00
F3	1380.00	1375.00	1377.50	295.67	293.33	294.50
F4	1510.00	1515.00	1500.50	345.33	350.66	348.00
F5	1375.00	1370.00	1372.50	312.33	310.33	311.30
F6	1485.00	1490.00	1440.00	322.33	325.67	324.00
Mean	1418.00	1419.00	1409.00	306.72	307.56	307.13
LSD at 5%	1 st season =2.51 2 nd season =2.13			1 st season =13.87 2 nd season =11.34		

F1= control 48 kg P₂O₅ (single super phosphate 16% P₂O₅).

F2= 48 kg P₂O₅ (phosphoric acid 85% P₂O₅)

F3= 48 kg P₂O₅ (24 kg P₂O₅ single super phosphate 16% P₂O₅ + 24 kg P₂O₅ phosphoric acid 85% P₂O₅).

F4= 48 kg P₂O₅ (mono ammonium phosphate MAP 12% N:61% P₂O₅).

F5= 48 kg P₂O₅ (24 kg P₂O₅ single super phosphate 16% P₂O₅ +24 kg P₂O₅ mono ammonium phosphate).

F6= 48 kg P₂O₅ (16 kg P₂O₅ single super phosphate 16% P₂O₅ + 16 kg P₂O₅ phosphoric acid 85% P₂O₅ +16 kg P₂O₅ mono ammonium phosphate MAP 12% N:61% P₂O₅).

These results coincided with those obtained by several authors such as Ragab *et al.*, (2015) on potatoes who found that the addition of phosphorus as MAP resulted in vigor plant growth expressed as plant height, number of leaves/plant, shoots/plant, fresh and dry weight, leaf area, leaf area index, relative growth rate and net assimilation rate if compared with triple superphosphate and diammonium phosphate (DAP). Maurya (1989) and Farahani *et al.*, (2009) found that the application of phosphorous fertilizer increased the vegetative yield of coriander. Also, these results are similar to the findings of

Bharose *et al.*, (2001) on Toria, Naderidarbaghshahi, *et al.*, (2011) on German chamomile (*Matricaria chamomilla* L.), Saharkhiz and Omidbaigi (2008) on feverfew, Sharafzadeh *et al.*, (2011) on sweet basil and Nassar *et al.*, (2004) on chamomile. They obtained the best results in higher levels of phosphorus.

These results may be due to increased available P concentrations in the plant's root zone. Because mono ammonium phosphate high-efficiency source of phosphorus as phosphate (PO_4^-) promotes rapid absorption by plants. It also has an acidifying effect on the favoring soiling solubility of phosphorus and micro-nutrients, fully soluble and fast dissolving. Although the amount of soil P concentration seems to be sufficient, the application of F4 48 kg P_2O_5 (mono ammonium phosphate), which produces the high available P-containing fertilizers in the soil, can increase the plant nutrient concentrations (Nell *et al.*, 2009; Rahimi *et al.*, 2013; Khalid 2012; Khalid, 2015; Kumar *et al.*, 2015).

3.1.2. Fresh and dry weights of flower heads (g/m^2)

Data presented in Table (5) indicated the main effect of fertilization on fresh and dry weights of flower heads (g/m^2). The highest values of these parameters were obtained when the plants were treated with F4 48 kg P_2O_5 (mono ammonium phosphate) compared to other fertilization treatments resulting in significant differences in most cases. The heaviest fresh weights of flower heads recorded 675 and 670 g/m^2 in the first and second seasons. It was followed by F6 48 kg P_2O_5 (16 kg P_2O_5 single super phosphate 16% P_2O_5 +16 kg P_2O_5 phosphoric acid 85% P_2O_5 +16 kg P_2O_5 mono ammonium phosphate MAP 12% N:61% P_2O_5) gave the lower values of fresh weights. Conversely, fertilization with control treatment F1 48 kg P_2O_5 (single super phosphate 16% P_2O_5) produced the lowest values of fresh weights of flower heads which recorded 675 and 670 g/m^2 in the first and second seasons, respectively. Concerning the effect of fertilization on the dry weight of flower heads, a significant increase was observed when the treatment of F4 was applied to plants in both seasons; they were 132.33 and 134.66 g/m^2 in the first and second seasons, respectively. It was followed by F6 48 kg P_2O_5 (16 kg P_2O_5 single super phosphate 16% P_2O_5 +16 kg P_2O_5 phosphoric acid 85% P_2O_5 +16 kg P_2O_5 mono ammonium phosphate MAP 12% N:61% P_2O_5) which recorded 131.33 and 130.33 g/m^2 in the first and second seasons, respectively. Meanwhile, control treatment F1 (48 kg P_2O_5 as single super phosphate 16% P_2O_5) produced the lowest dry weights of flower heads, which were 115.33 and 116.33 g/m^2 in the first and second seasons, respectively. The differences between treatments and control were significant when the experiment was repeated in both seasons.

Table 5: Effect of different phosphorus fertilization resources on the fresh and dry weights of flower heads (g/m^2) during 2018/2019 and 2019/2020 seasons

Treatments	Fresh weight of flower heads (g/m^2)			Dry weight of flower heads (g/m^2)		
	1 st season	2 nd season	Mean	1 st season	2 nd season	Mean
F1	570.00	580.00	575.00	115.33	116.33	115.83
F2	610.00	600.00	605.00	129.33	130.33	129.83
F3	645.00	635.00	640.00	124.66	127.66	126.16
F4	675.00	670.00	672.50	132.33	134.66	133.50
F5	650.00	655.00	652.50	125.33	123.66	124.50
F6	670.00	665.00	662.50	131.33	130.33	130.83
Mean	636.67	634.17	634.58	126.39	127.16	126.78
LSD at 5%	1 st season =4.36 2 nd season =3.03			1 st season =3.51 2 nd season =3.33		

F1= control 48 kg P_2O_5 (single super phosphate 16% P_2O_5).

F2= 48 kg P_2O_5 (phosphoric acid 85% P_2O_5) .

F3= 48 kg P_2O_5 (24 kg P_2O_5 single super phosphate 16% P_2O_5 + 24 kg P_2O_5 phosphoric acid 85% P_2O_5).

F4= 48 kg P_2O_5 (mono ammonium phosphate MAP 12% N:61% P_2O_5).

F5= 48 kg P_2O_5 (24 kg P_2O_5 single super phosphate 16% P_2O_5 +24 kg P_2O_5 mono ammonium phosphate).

F6= 48 kg P_2O_5 (16 kg P_2O_5 single super phosphate 16% P_2O_5 + 16 kg P_2O_5 phosphoric acid 85% P_2O_5 +16 kg P_2O_5 mono ammonium phosphate MAP 12% N:61% P_2O_5).

Our results agree with those obtained by Maurya (1989) and Farahani *et al.*, (2009) findings that the application of phosphorous fertilizer increased the vegetative yield of coriander. Also, these results

are similar to the findings of Bharose *et al.*, (2001) on toria, Naderidarbaghshahi *et al.*, (2011) on German chamomile, Saharkhiz and Omidbaigi (2008) on feverfew and Sharafzadeh *et al.*, (2011) on sweet basil also Nassar *et al.*, (2004) on chamomile obtained the best results in higher levels of phosphorus. Generally, it could be concluded that fresh and dry weights of flower heads (g/m²) recorded their maximum values when phosphorus was added as a MAP source. The opposite was true for single super phosphate treatment. The highest chamomile yield was obtained by using MAP. It may be due to enhanced development of meristematic tissue or the increment of the following attributes: the number of flower heads, photosynthetic activity, mobilization of photosynthates, and chlorophyll biosynthesis photosynthetic of CO₂ fixation, cell division, and carbohydrates metabolism. Also, it could be concluded that the superiority of MAP may be attributed to its higher water solubility and pH value, and higher nitrogen content. These results agree with Hill *et al.*, (2011) and Rosen *et al.*, (2014).

3.2. Chemical constituents

3.2.1. Essential oil percentage

Data in Table (6) showed a progressive increase in the volatile oil percentage of the chamomile plant during both seasons. The differences between fertilization treatments were significant. The highest oil percentage (%) in flower heads was produced when the plants were treated with F4 (48 kg P₂O₅ mono ammonium phosphate) compared with other fertilization treatments; these estimates were 0.77 and 0.75% in the first and second seasons, respectively. It was followed by F6 48 kg P₂O₅ (16 kg P₂O₅ single super phosphate 16% P₂O₅) + 16 kg P₂O₅ phosphoric acid 85% P₂O₅ + 16 kg P₂O₅ mono ammonium phosphate MAP 12% N:61% P₂O₅) which gave the following higher values of oil percentage (%) which were recorded 0.75 and 0.74% in the first and second season, respectively. In contrast, control treatment F1 48 kg P₂O₅ (single super phosphate 16% P₂O₅) produced the lowest oil percentage (%) in flower heads during both seasons.

Table 6: Effect of different phosphorus fertilization resources on essential oil percentage in flower heads and oil yield per m² during 2018/2019 and 2019/2020 seasons

Treatments	Essential oil percentage (%) in flower heads			Oil yield per (ml/m ²) in flower heads		
	1 st season	2 nd season	Mean	1 st season	2 nd season	Mean
F1	0.70	0.72	0.71	0.81	0.84	0.83
F2	0.73	0.74	0.74	0.94	0.96	0.95
F3	0.73	0.73	0.73	0.91	0.93	0.92
F4	0.77	0.75	0.76	1.02	1.01	1.02
F5	0.73	0.75	0.74	0.91	0.93	0.92
F6	0.75	0.74	0.75	0.98	0.96	0.97
Mean	0.74	0.73	0.74	0.94	0.93	0.94
LSD at 5%	1 st season = 0.01 2 nd season = 0.01			1 st season = 4.36 2 nd season = 3.03		

F1= control 48 kg P₂O₅ (single super phosphate 16% P₂O₅).

F2= 48 kg P₂O₅ (phosphoric acid 85% P₂O₅) .

F3= 48 kg P₂O₅ (24 kg P₂O₅ single super phosphate 16% P₂O₅ + 24 kg P₂O₅ phosphoric acid 85% P₂O₅).

F4= 48 kg P₂O₅ (mono ammonium phosphate MAP 12% N:61% P₂O₅).

F5= 48 kg P₂O₅ (24 kg P₂O₅ single super phosphate 16% P₂O₅ + 24 kg P₂O₅ mono ammonium phosphate).

F6= 48 kg P₂O₅ (16 kg P₂O₅ single super phosphate 16% P₂O₅ + 16 kg P₂O₅ phosphoric acid 85% P₂O₅ + 16 kg P₂O₅ mono ammonium phosphate MAP 12% N:61% P₂O₅).

Effect of different phosphorus fertilizers on essential oil percentage may be due to their distinct effect on enzyme activity and metabolism necessary for oil production. According to Jeshni *et al.*, (2015), P fertilization increased the essential oil concentration of chamomile. Khalid (1996) and Economakis (2002) showed that P significantly increased the essential oil content of Apiaceae and *Origanum dictamnus* plants. P enhanced essential oil extracted from bitter fennel (*Foeniculum vulgare* Mill.) (Tank *et al.*, 2007; Osman 2009). P had a significant effect on the essential oil content isolated from sweet fennel cultivars growing in arid regions in Egypt (Zaki *et al.*, 2010; 2011). Also, it increased the essential oil content of basil (Ramezani *et al.*, 2009), fennel (*Foeniculum vulgare*) (Kapoor *et al.*,

2004), and cumin (Tuncturk and Tuncturk, 2006). Furthermore, a lack of effect the phosphorous application on oil percentage was reported for other plants, including *Salvia officinalis* (Nell *et al.*, 2009; Rioba *et al.*, 2015).

3.2.2. Essential oil yield (ml/m²)

Data of the volatile oil yield productivity are shown in table (6). The results showed that the best effect in this respect was obtained by fertilized chamomile plant with F4 showed its superiority for increasing the oil yield per (ml/m²) as it gave 1.02 and 1.01 ml/m² followed descendingly by using the treatment F6, which recorded 0.98 and 0.96 ml/m² in the first and second seasons, respectively. Conversely, the minor oil yield per plant (ml/m²) was obtained by treating the plants with F1, recorded at 0.81 and 0.84 ml/m². This may be due to this treatment gave less plant growth, reflecting less oil yield per plant. The increment of oil yield per plant may be due to the beneficial effects of these fertilizers on the synthesis of carbohydrates in plant tissues, consequently used for oil production. The obtained results follow those Al-Humaid (2004) reported on fennel plants and Khalid (2015) on coriander. They stated that the application of phosphorous caused a pronounced increment in essential oil yield.

3.2.3. Essential oil constituents

Six main constituents were identified in the essential oil extracted from chamomile flowers head, accounting for 79.15 - 86.31% of total components (Table 7). The main constituents of chamomile flowers' essential oil as detected by GC/MS were chamazulene, bisabolol oxide B, and α -bisabolol, which increased by F4 48 kg P₂O₅ (mono ammonium phosphate MAP 12% N:61% P₂O₅) treatment compared with the control treatment F1 48 kg P₂O₅ (single super phosphate 16% P₂O₅). Also, the chemical classes of chamomile essential oil were changed in all treatments compared with the control. It could be concluded that the superiority of F4 may be attributed to its high effect on enzymes activity and accelerate plant metabolism.

The previous investigations concerning the response of chamomile yield and its components to the various phosphorus sources were confirmed (Sheibanivaziri, 1997; Rosen and McNearney, 2003; Hill *et al.*, 2011; Rosen *et al.*, 2014; Nikolova *et al.*, 1999). They showed that P fertilization increased the essential oil concentration of chamomile, and it is a necessary element. Thus, the essential oil was increased by applied MAP as expected in the present study. Phosphorus also has various cellular functions in plants, including signaling and transmembrane metabolic flux. The results were in line with those obtained by Marschner (1999) and Nilbe *et al.*, (2005), who observed increasing biomass of chamomile. Therefore, secondary metabolism (essential oil) is modulated by these mechanisms.

Table 7: Effect of different phosphorus fertilization resources on the main constituents of essential oil during 2018/2019 and 2019/2020 seasons

Treatments	Germacrene D	α -Farnesene	Bisabolol Oxide B	α -Bisabolol	Chamazulene	Bisabolol oxide A	Total identified
F1	4.75	6.47	12.85	12.76	41.52	6.76	82.10
F2	4.54	7.32	12.73	10.83	41.66	6.35	79.15
F3	4.43	7.61	13.52	11.51	40.88	6.72	81.83
F4	4.91	7.55	13.61	11.65	41.68	7.85	86.31
F5	4.88	6.92	12.60	10.89	41.44	6.89	83.56
F6	4.74	7.32	12.89	11.32	41.35	7.71	80.48

F1= control 48 kg P₂O₅ (single super phosphate 16% P₂O₅).

F2= 48 kg P₂O₅ (phosphoric acid 85% P₂O₅).

F3= 48 kg P₂O₅ (24 kg P₂O₅ single super phosphate 16% P₂O₅ + 24 kg P₂O₅ phosphoric acid 85% P₂O₅).

F4= 48 kg P₂O₅ (mono ammonium phosphate MAP 12% N:61% P₂O₅),

F5= 48 kg P₂O₅ (24 kg P₂O₅ single super phosphate 16% P₂O₅ +24 kg P₂O₅ mono ammonium phosphate).

F6= 48 kg P₂O₅ (16 kg P₂O₅ single super phosphate 16% P₂O₅ + 16 kg P₂O₅ phosphoric acid 85% P₂O₅ +16 kg P₂O₅ mono ammonium phosphate MAP 12% N:61% P₂O₅).

3.2.4. Mineral elements content (NPK)

Data presented in Table (8) showed that fertilized chamomile plants with F4 48 kg P₂O₅ (mono ammonium phosphate) treatment showed its superiority for increasing nitrogen, phosphorus, and potassium percentages in their tissues. It was followed descendingly by using F6 compared with the other phosphorus fertilizers. The positive effect of F4 48 kg P₂O₅ (mono ammonium phosphate) on mineral composition could be attributed to the high available P-containing fertilizers in the soil, increasing the plant nutrient concentrations. It may be due to its translocation from the root to the above-ground parts and more absorption and accumulation of N, P, and K. The obtained results for the effect of P on the percentages of the mineral are in harmony with those mentioned by (Massoud, 2006; El-Sayed *et al.*, 2011; Abdel-Razzak *et al.*, 2013; Rosen *et al.*, 2014).

Table 8: Effect of different phosphorus fertilization resources on nitrogen, phosphorus and potassium (%) during 2018/2019 and 2019/2020 seasons

Treatments	N			P			K		
	1 st season	2 nd season	Mean	1 st season	2 nd season	Mean	1 st season	2 nd season	Mean
F1	1.81	1.83	1.82	0.35	0.34	0.35	2.11	2.13	2.12
F2	2.31	2.29	2.30	0.43	0.44	0.44	2.29	2.27	2.28
F3	2.22	2.20	2.21	0.41	0.41	0.41	2.24	2.25	2.25
F4	2.35	2.36	2.36	0.47	0.48	0.48	2.31	2.33	2.32
F5	2.11	2.12	2.12	0.42	0.43	0.43	2.14	2.16	2.15
F6	2.25	2.25	2.25	0.43	0.44	0.44	2.15	2.17	2.16
LSD at 5%	1 st season = 0.08 2 nd season = 0.09			1 st season = 0.02 2 nd season = 0.02			1 st season = 0.06 2 nd season = 0.07		

F1= control 48 kg P₂O₅ (single super phosphate 16% P₂O₅).

F2= 48 kg P₂O₅ (phosphoric acid 85% P₂O₅)

F3= 48 kg P₂O₅ (24 kg P₂O₅ single super phosphate 16% P₂O₅ + 24 kg P₂O₅ phosphoric acid 85% P₂O₅).

F4= 48 kg P₂O₅ (mono ammonium phosphate MAP 12% N:61% P₂O₅).

F5= 48 kg P₂O₅ (24 kg P₂O₅ single super phosphate 16% P₂O₅ + 24 kg P₂O₅ mono ammonium phosphate).

F6= 48 kg P₂O₅ (16 kg P₂O₅ single super phosphate 16% P₂O₅ + 16 kg P₂O₅ phosphoric acid 85% P₂O₅ + 16 kg P₂O₅ mono ammonium phosphate MAP 12% N:61% P₂O₅).

Finally, it could be concluded that the highest plant growth vigor was obtained from using F4 (MAP) followed by F6 48 kg P₂O₅ (16 kg P₂O₅ single super phosphate 16% P₂O₅ + 16 kg P₂O₅ phosphoric acid 85% P₂O₅ + 16 kg P₂O₅ mono ammonium phosphate MAP 12% N:61% P₂O₅). The increase in dry weights of flowers head may be attributed to the rise in fresh plant weight, leading to more active photosynthesis and, in turn, more carbohydrate accumulation. The higher values of essential oil in flower heads may be due to the improvement effect of F4 (MAP) and the treated plants by F6 on fresh weight of flower heads and the advancement effect on essential oil in flower heads or oil yield per ml/m². The advantages of MAP as a phosphorus source on the productivity of chamomile plants might be attributed to one or more of the following. The higher nitrogen content (12 %), the higher phosphorus content (61%), water solubility (370 g/l), and its pH value (4.0 to 4.5), compared by 5 ppm max Nitrate (NO₃) and 85% P₂O₅, 548 g/100 ml (20°C), solubility and 2.14 pH value for phosphoric acid and pH value 1.5 (20°C, 10 g/l). Because of properties variation within MAP, phosphoric acid, and single super phosphate, results clearly showed that the differences were significant concerning most plant growth parameters. The obtained results are in good accordance with those reported by Balemi and Schenk (2009), and Rosen *et al.*, (2014).

4. Conclusion

The experimental results showed that applying F4 48 kg P₂O₅/feddan (mono ammonium phosphate) produced the highest plant growth characteristics, oil content, N, P, and K in plant tissues. Also, this treatment increased the concentrations of chamazulene to (41.68 %) and bisabolol oxide B to (13.61%). In comparison, super-phosphate as phosphorus fertilizer gave the lowest values in the two tested seasons.

References

- Abd El-Aleem, W.H., E.S. Hamed and W.I.M. Toaima, 2021. Effect of blue green algae extract on three different curly parsley varieties under Sinai conditions. *Bulgarian Journal of Agricultural Science*, 27 (5): 887-895.
- Abdel-Razzak, H.S., A.G. Moussa, M.A. Abd El-Fattah and G.A. El-Morabet, 2013. Response of sweet potato to integrated effect of chemical and natural phosphorus fertilizer and their levels in combination with mycorrhizal inoculation. *Journal of Biological Sciences*, 13: 112-122.
- Abla, H., M.F. Nassar, N.S. Hashim and H. Abo-zaid, 2004. Effect of gamma irradiation and phosphorus on growth and oil production of chamomile (*Chamomilla recutita* L. Rauschert). *Intl. J. Agric. Biol.* 6(5): 776-780.
- Adams, R.P., 2007. Identification of Essential Oil Components by Gas Chromatography/Quadrupole Mass Spectroscopy, 4th ed., Allured Publishing Corporation, Illinois.
- Ahmadian, A., A. Ghanbari, B. Siahshar, M. Haydari, M. Ramroodi and S.M. Mousavinik, 2011. Study of chamomile's yield and its components under drought stress and organic and inorganic fertilizers using and their residue. *J Micro. Antimicrob.*, 3(2): 23-28.
- Al-Humaid, A.L., 2004. Effect of bio and chemical fertilizers on the growth and essential oil yield of fennel (*Foeniculum vulgare*, Mill.) Alex. *J Agric. Res.*, 49 (3):83-91.
- Babushok, V.I., P.J. Linstrom and I.G. Zenkevich, 2011. Retention indices for frequently reported compounds of plant essential oils. *Journal of Physical and Chemical Reference Data* 40, 043101-47.
- Balemi, T. and M.K. Schenk, 2009. Genotypic variation of potato for phosphorus efficiency and quantification of phosphorus uptake with respect to root characteristics. *Journal of Plant Nutrition and Soil Science*, 172(5): 669-677.
- Bharose, R., S. Chandra, T. Thomas and D. Dhan, 2001. Effect of different levels of phosphorus and sulphur on yield and availability of N P K, protein and oil content in Toria (Brassica sp.) var. p.t.-303. *ARPN J. Agric. and Biol. Sci.* 6(2): 31-33.
- British Pharmacopoeia, 1963. Determination of Volatile Oil in Drugs, published by the Pharmaceutical Press, London.
- Brown, J.D. and D. Lilleland, 1946. Rapid determination of potassium and sodium in plant material and soil extracts by flame photometry. *Proc. Amer. Soc. Hort. Sci.* 48: 341-346.
- Bunzen, J.N.P., Guichard, J. Labbr and J. Trenchant, 1969. In "Practical Manual of Gas Chromatography". (J. Trenchant, ed.), El-Seivier, Publ. Comp., Amesterdam, Netherland.
- Chapman, H.D. and P.F. Pratt, 1978. Methods of Analysis for Soils, Plants and Water. Univ. of California, Prical Publication, 4030, 12-19.
- Das, M., 2014. Chamomile: Medicinal, Biochemical, and Agricultural Aspects. CRC Press, 316 pp.
- Economakis, G., H. Skaltsa, G. Demetzos, M. Sokovic and G.A. Thanos, 2002. Effect of phosphorus concentration of the nutrient solution on the volatile constituents of leaves and bracts of (*Origanum dictamnus*). *Journal of Agriculture and Food Chemistry*, 50 (22): 6276-6280.
- El-Sayed, H.E.A., A.S. El-Dean, S. Ezzat and A.H.A. El-Morsy, 2011. Responses of productivity and quality of sweet potato to phosphorus fertilizer rates and application methods of the humic acid. *Int. Res. J.Agric. Sci. Soil Sci.*, 1: 383-393.
- Essa, A.M., 1999. Effect of some cultural treatments on growth and yield of (*Marticaria chamomilla* L.) plant, Ph.D. Thesis, Fac. Agric. Ain Shams Univ. Cairo, Egypt.
- Farahani, H.A., S.A. Valadabadi and M.A. Khalvati, 2009. Interactive effects of P supply and drought on root growth of the mycorrhizal coriander (*Coriandrum sativum* L.). *J. Plant Breeding and Crop Sci.* 1(5): 217-222.
- Hamed, E.S., 2018. Effect of biofertilization on Siwa Oasis mint (*Mentha spicata* L. cv. Siwa) plants. *Journal of Medicinal Plants Studies*, 6 (3): 24-33.
- Hill, M.W., B.G. Hopkins. C.J. Ransom, and B.L. Webb., 2011. Improving phosphorus use efficiency with Carbond P. In: Proceedings of the Western Nutrient Management Conference, 9: 129-134. Reno, NV, USA: CSREES WERA-103 Nutrient.
- Hoftman, E., 1967. Chromotography, Reinhild. Corp., 2nd Ed., 208-515. 131
- Jennings, W. and T. Shibamoto, 1980. Qualitative analysis of flavor and fragrance volatiles by Glass capillary gas chromatograph, 1st ed., Academic Press, New York.

- Jeshni, M.G., M. Mousavinik, I. Khammari and M. Rahimi, 2015. The changes of yield and essential oil components of 5 German Chamomile (*Matricaria recutita* L.) under 6 application of phosphorus and zinc fertilizers and 7 drought stress conditions. *Journal of the Saudi Society of Agricultural Sciences* 16(1): 60-65.
- Kandeel, A.M.A., 1982. Effect of some agricultural treatments on the yield and oil percentage of chamomile plant *Marticaria chamomilla* L, M.Sc. Thesis, Fac. Agric. Ain Shams Univ., Cairo, Egypt.
- Kapoor, R., B. Giri and K.G. Mukerji, 2004. Improved growth and essential oil yield and quality in (*Foeniculum vulgare* Mill). on mycorrhizal inoculation supplemented with P-fertilizer. *Bioresource Technology*, 93: 307-311.
- Khalid, K.A., 2006. Influence of water stress on growth, essential oil, and chemical composition of herbs, *Ocimum* sp. *Int. Agrophys*, 20: 289-296.
- Khalid, K.A., 2012. Effect of phosphorous fertilization on anise, coriander and sweet fennel plants growing under arid region conditions. *Med. Aromat. Plant. Sci. Biotechnol.* 6(1):127-131.
- Khalid, K.A., 2015. Effect of macro and micro nutrients on essential oil of coriander fruits. *JMES* 6(8):2060-2065.
- Khalid, K.A., 1996. Effect of fertilization on the growth, yield and chemical composition of some medicinal umbelleferous plant. M.Sc. Thesis, Faculty of Agriculture, Al-Azhar University, Cairo, Egypt.
- Koch, F.C. and T.L. Mc-Meekin, 1924. A new direct nesslerization micro-Kjeldahl method and modification of the Nessler folia reagent for ammonia. *J. Amer. Chem. Soc.* 46: 2066.
- Kumar, P., S.K. Pandey and P. Kumar, 2015. Effect of different phosphorus levels on nutrient content, uptake and economics of urdbean under custard apple based agri-horti system. *J. Agri Search*, 2(2):88-93.
- Letchamo, W., A. Gosselin and G. Lisin, 2006. Chamomile varieties and quality improvement issues. Program and Abstract Book of the 1th. International Symposium on Chamomile Research, Development and Production. Presov University in Presov.
- Marschner, H., 1999. Mineral Nutrition of Higher Plants, Academic Press, London.
- Massada, Y., 1976. Analysis of Essential Oils by Gas Chromatography and Mass Spectrometry, Illustrated, ed. Wiley & Sons, New York.
- Massoud, Hekmat Y.A., 2006. Effect of phosphorus fertilization levels and foliar application with active dry yeast bio-fertilizer on growth, herb yield, essential oil productivity and chemical components of sage (*Salvia officinalis*, L.). *Journal of Plant Production*, 31 (10): 6649-6665.
- Maurya, K.R., 1989. Growth, yield and quality component in coriander genotypes. *Indian J. Hort.* 46(1): 107-110.
- Naderidarbaghshahi, M., S.M. Monemian, H. Zeynali and B. Bahari, 2011. Effects of different levels of nitrogen, phosphorus and potassium fertilizers on some agro-morphological and biochemical traits of German chamomile (*Matricaria chamomilla* L.). *J. Medicinal Plants Res.* 6(2): 277-283.
- Nassar, A.H., M.F. Hashim, N.S. Hassan and H. Abo-Zaid, 2004. Effect of gamma irradiation and phosphorus on growth and oil production of chamomile (*Chamomilla recutita* L. Rauschert). *Int. J. Agri. Biol.* 6(5): 776-780.
- National Committee for Clinical Laboratory Standards, 2002. Performance Standards for Antimicrobial Susceptibility Testing. 29th ed., Clinical and Laboratory Standards Institute, Wayne.
- Nell, Monika, *et al.*, 2009. Effect of phosphorus uptake on growth and secondary metabolites of garden sage (*Salvia officinalis* L.). *Journal of the Science of Food and Agriculture*, 89 (6): 1090-1096.
- Nikolova, A., A.K. Kozuharova, V.D. Zheljzakov and L.E. Craker, 1999. Mineral nutrition of chamomile (*Chamomilla recutita* (L.). *Acta Hort.* 502 203.
- Nilbe, C.M., C.M. Vieira, A. Néstor, Z. Heredia and J.M. Siqueira, 2005. Biomass production of chamomile capitula as a result of nitrogen and phosphorus. *Hort Braz* 23(1): 32-37.
- Omidbeigi R., 2005. Pharmaceutical Plants Production, Fekr-e-Rooz press, Iran.
- Osman, Y.A.H., 2009. Comparative study of some agricultural treatments effects on plant growth, yield and chemical constituents of some fennel varieties under Sinai conditions. *Research Journal of Agriculture and Biological Sciences* 5(4):541-554.

- Osman, Y.A.H., M.A. El-Wahab and F.A. Ali, 2013. Production of Aromatic Flowering and Fruiting and Roots Plants in New Reclaimed Land. Pub. Ministry of Agriculture, Cairo, ISBN: 978-977-302-378-2 [In Arabic].
- Presibella, M.M., L.B. Villas-Bôas, M.S. Belletti, C.A.M. Santos and A.M. WeffortSantos, 2006. Comparison of chemical constituents of *Chamomilla recutita* L. Rauschert essential oil and its antichemotactic activity. *Brazilian Archiv. Biol. Technol.* 49(5): 717-724.
- Ragab, M.E., M.M. Soliman, F.A. Rizk and S.H. Mahmoud, 2015. Effect of different phosphorus sources on the plant growth, tubers yield and nutritional value of potatoes. *Middle East Journal of Agric. Research.* 4(2): 388-394.
- Rahimi, M.F., F. Reza, and M. Rahman 2013. Effects of potassium and phosphorus fertilizers on arsenic phytotoxicity and essential oil yield of two basil cultivars. *BEPLS* 2(5):86-92.
- Ramezani S., Z.A.A. Aziz and S.Z.A. Idid, 2009. Public space and conservation of a historic living city: Melaka, Malaysia. Retrieved May 20, 2014 from <http://www.epublication.fab.utm.my/252/1/routledgevol42010.pdf>
- Rioba, N.B., F.M. Itulya, M. Saidi, N. Dudai and N. Bernstein, 2015. Effects of nitrogen, phosphorus and irrigation frequency on essential oil content and composition of sage (*Salvia officinalis* L.). *Journal of Applied Research on Medicinal and Aromatic Plants* 2:21-29.
- Rosen, C.J., A.K. Kelling, J.C. Stark and A.P. Gregory, 2014. Optimizing phosphorus fertilizer management in potato production. *Am. J. Potato Res* 91:145–160.
- Rosen, C.R. and M. McNearney, 2003. Potato yield and tuber set as affected by phosphorus fertilization. *Field Research in Soil Science*, 34-38.
- Saharkhiz, M.J. and R. Omidbaigi, 2008. The effect of phosphorus on the productivity of feverfew (*Tanacetum parthenium* (L.) Schultz Bip). *Advances in Natural and Applied Sci.* 2(2): 63-67.
- Salmon, I. and E.N. Abou-zied, 2006. The qualitative-quantitative characteristics of chamomile essential oil in Egypt. Program and Abstract book of the 1st International Symposium on Chamomile Research, Development and Production, 41.
- Sharafzadeh, S., M. Esmaeili and A.H. Mohammadi, 2011. Interaction effects of nitrogen, phosphorus and potassium on growth, essence and total Phenolic content of Sweet basil. *Advances in Environmental Biology*, 5(6): 1285-1289.
- Sheibanivaziri, M., 1997. Effect of nitrogen, phosphorus and potassium fertilizers on essence and chamazulen amount of chamomile flowers, PhD dissertation, Isfahan Medical University, Iran.
- Snedecor, G.W. and W.G. Cochran, 1968. *Statistical Methods* sixth edition, The Iowa State Univ. Press, Ames, Iowa, U.S.A
- Tank, D.A., T.G. Meisheri and V.P. Usadadia, 2007. Effects of spacing, nitrogen and phosphorus on growth and yield of fennel (*Foeniculum vulgare* Mill.). *Research Grops* 7(2): 561-562.
- Toaima, W.I.M., M.Y.M.A. Badawy and E.S. Hamed, 2022. Effect of organic fertilization on productivity of some newly introduced basil varieties under Siwa Oasis conditions. *Journal of Applied Biology & Biotechnology*, 10 (02): 74-88.
- Troug, E. and A. Mayer, 1939. Improvement in the direness colorimetric method for phosphorus and arsenic. *Ind. Eng. Chem. Anal. Ed.*, 1:136-139.
- Tuncturk, R. and M. Tuncturk, 2006. Effects of different phosphorus levels on the yield and quality components of cumin (*Cuminum cyminum* L.). *Res. J. Agric. Biol. Sci.* 2: 336-340.
- Zaki, M.F., A.A. Abdelhafez and C.Y. El-Dewiny, 2010. Influence of applying phosphate bio-fertilizers and different levels of phosphorus sources on the productivity, quality and chemical composition of sweet fennel (*Foeniculum vulgare* Mill.). *Australian Journal of Basic and Applied Science*, 4(2):334-347.
- Zaki, M.F., H.A.M. Habib, E.H. Abdelsamad and Z.A. Lashine, 2011. Effect of different sources and rates of phosphorus on vegetative growth, antioxidant and mineral contents of two sweet fennel cultivars. *European Journal of Scientific Research*, 4:499-511.